

The non-abelian squares are not context-free

Shuo Tan

School of Computer Science

University of Waterloo

Waterloo, ON N2L 3G1

Canada

sh22tan@gmail.com

October 20, 2011

Erdős introduced the notion of *abelian square* in a 1961 paper [1]:

Definition 1. A word w is an *abelian square* if there exist two words s and t such that $w = st$ and t is a permutation of s .

At the DLT 2011 conference in Milan, Italy, Maxime Crochemore asked if the language of non-abelian squares is context-free. In this note, we answer his question.

Let $\Sigma = \{0, 1\}$ be the alphabet in this context. Let L denote the language of all words that are not abelian squares over Σ . Our goal is to prove the following:

Theorem 2. L is not context-free.

In order to prove this, we introduce some notation. Define $w_i = 10^{i-1}$ for all $i > 1$. Let $R = w_4^* w_3 w_2^* w_3 w_3^*$.

Lemma 3. The word $w_4^n w_3 w_2^{n!+n} w_3 w_3^{2(n!+n)} \in L \cap (\Sigma^2)^* \cap R$.

Proof. Suppose $z = w_4^n w_3 w_2^{n!+n} w_3 w_3^{2(n!+n)}$. Clearly $z \in (\Sigma^2)^* \cap R$. Suppose m is the total number of 1's in z . Then $m = 3n! + 4n + 2$. However, the number of 1's in the second half of z is $\frac{4n!}{3} + 2n + 1$, which is not equal to $\frac{m}{2}$. Hence z is not an abelian square. Thus $z \in L$. $\square$

Definition 4. Let $w = 0^{s_0} 10^{s_1} \dots 10^{s_k}$ be a word over the alphabet Σ . We define

$$\text{alt}(w) = |\{1 \leq i < k : s_i \neq s_{i+1}\}|.$$

Define $\text{alt}(\cdot)$ over a language K as follows: $\text{alt}(K) = \max_{w \in K} \text{alt}(w)$.

Definition 5. A sequence of non-negative integers $(a_k)_{k=1}^n$ is called *uneven* if $n > 1$ and $\exists i \in [1, n]$ such that $a_i \neq a_{i+1}$. Here $a_{n+1} = a_1$. Otherwise, it is called *even*.

Definition 6. Suppose w is a word that contains a 1. Then w is called *uneven* if the sequence $(s_1 + s_{k+1}, s_2, \dots, s_k)$ is uneven, where $w = 0^{s_1}10^{s_2}1 \dots 0^{s_k}10^{s_{k+1}}$. Otherwise, it is called *even*.

Example 7. Suppose w is an even word and of the form $0^{s_1}10^{s_2}1 \dots 0^{s_k}10^{s_{k+1}}$. Now we consider what w looks like. Since w is even, we get that $s_1 + s_{k+1} = s_2 = \dots = s_k$. Then $w = 0^s 1 (0^{s+t} 1)^k 0^t$ for some $s, t, k \geq 0$.

Lemma 8. If w is an uneven word, then $\text{alt}(w^k) \geq k - 1$.

Proof. Suppose $w = 0^{s_1}10^{s_2}1 \dots 0^{s_k}10^{s_{k+1}}$. Then $w^k = 0^{s_1}(10^{s_2} \dots 10^{s_k}10^{s_{k+1}+s_1})^{k-1}10^{s_2} \dots 10^{s_k}10^{s_{k+1}}$. Since w is uneven, we get $\text{alt}(10^{s_2} \dots 10^{s_k}10^{s_{k+1}+s_1}) \geq 1$. It follows that $\text{alt}(w^k) \geq k - 1$. $\square$

Lemma 9. Let $T = L \cap (\Sigma^2)^* \cap R$. Then T is not context-free.

Proof. (By Ogden's Lemma) For any $n > 4$, let $z = w_4^n w_3 w_2^{n!+n} w_3 w_3^{2(n!+n)}$. By Lemma 3 we see that $z \in T$. Mark the first $4n$ bits of z , that is, the bits corresponding to w_4^n . Let $m(s)$ denote the number of bits marked in s . Now we show by contradiction that no decomposition $z = u_0 v_0 w_0 x_0 y_0$ satisfies all the following three conditions:

1. condition A: $m(v_0 x_0) > 0$
2. condition B: $m(v_0 w_0 x_0) \leq n$
3. condition C: $\forall i \geq 0, u_0 v_0^i w_0 x_0^i y_0 \in T$

Before further consideration into all possible decompositions, we first mark z with different colors. Mark the bits corresponding to w_4^n red. Mark the next 3 bits corresponding to w_3 blue. Mark the bits corresponding to $w_2^{n!+n}$ green. Mark the bits corresponding to $w_3 w_3^{2(n!+n)}$ black. Define a new function $m(\text{color}, x)$ as the number of bits in x colored color. Note that $m(x)$ in our former definition is the same as $m(\text{red}, x)$. Here is a picture of how z is colored:

$$\underbrace{w_4 w_4 \dots w_4}_{\text{red}} \underbrace{w_3}_{\text{blue}} \underbrace{w_2 \dots w_2}_{\text{green}} \underbrace{w_3 w_3 \dots w_3}_{\text{black}}$$

Now we list all possible cases.

- (i) Either v or x is the empty word. Without loss of generality, suppose x is empty.
 - (i) $v \in 0^+$.
 - (ii) v contains a 1 and v is uneven.
 - (iii) v contains a 1 and v is even.
- (ii) Both v and x are non-empty words.
 - (i) $v \in 0^+$ or $x \in 0^+$.

- (ii) Both v and x contain a 1; v is uneven or x is uneven.
- (iii) Both v and x contain a 1 and are even.
 - (i) $m(\text{red}, v) = 0$, which means that w_4^n precedes v in z .
 - (ii) $m(\text{red}, v) > 0$.
 - (i) $m(\text{red}, x) > 0$.
 - (ii) $m(\text{red}, x) = 0$ and $m(\text{green}, x) > 0$
 - $m(\text{blue}, x) = 3$.
 - $m(\text{blue}, x) = 2$.
 - $m(\text{blue}, x) = 1$.
 - $m(\text{blue}, x) = 0$ and $m(\text{green}, x) > 1$.
 - $m(\text{blue}, x) = 0$ and $m(\text{green}, x) = 1$.
 - (iii) $m(\text{red}, x) = m(\text{green}, x) = 0$ and $m(\text{black}, x) > 0$

Suppose there exists a decomposition $z = uvwxy$ satisfying the above three conditions simultaneously.

Case i: First we consider the case when either v or x is empty. Without loss of generality, suppose x is empty. Then v cannot be empty, since vx is non-empty.

Case i.i: Suppose $v = 0^k$ for some $k \in \mathbb{N}^+$. Then we select $i = 4$. Since there are more than 3 successive 0's in v^4 , this is also true for uv^4wx^4y . However, no word in T contains more than 3 successive 0's. Hence we get a contradiction.

Case i.ii: Suppose v contains a 1 and is uneven. We pick $i = 6$. Then $\text{alt}(v^6) \geq 5 > \text{alt}(R) = 4$ by Lemma 8. So $uv^6wx^6y \notin T$, which violates condition C.

Case i.iii: Now we consider when v is even. In this case v can be written in the form $0^k 1 (0^{k+s} 1)^p 0^s$ for some $k, s, p \in \mathbb{N}$ (as we mentioned in Example 7). Then it follows that $m(\text{green}, v) = 0$ and $k + s = 3$ by the following argument. Suppose $m(\text{green}, v) > 0$. Then $m(\text{blue}, v) = 3$. That is to say, the w_3 between w_4 's and w_2 's lies in v . Then v must be of the form $r_1 01001 r_2$ for some words r_1 and r_2 . It follows that $k + s = 2$, since v is even. Now we select $i = 2$. Then $uv^2wx^2y = w_4^n w_3^l w_2^{n!+n} w_3 w_3^{2(n!+n)}$ for some $l > 1$, which violates condition C. Now suppose $k + s \neq 3$. Then we pick $i = 2$. It follows that uv^2wx^2y is of the form $w_4^l w_{k+s+1}^{2+2p} w_4^j w_3 w_2^{n!+n} w_3 w_3^{2(n!+n)} \notin T$, which violates condition C again. Now let $i = \frac{n!}{1+p}$. It follows that $z_i = uv^iwx^iy = w_4^{n!+n} w_3 w_2^{n!+n} w_3 w_3^{2(n!+n)} = (w_4^{n!+n} w_3 w_2^{n!+n})(w_3 w_3^{2(n!+n)})$ is an abelian square, a contradiction.

Case ii: Both v and x are non-empty. In this case, we first show that both v and x contain a 1. Then, we show v and x are even. Finally we rule out all subcases under the condition that v and x are even.

Case ii.i: Suppose $v = 0^k$ or $x = 0^l$ for some $k, l \in \mathbb{N}^+$. By a similar analysis in Case i.i, we get that this case violates condition C.

Case ii.ii: Suppose v is uneven. By a similar analysis in Case i.ii, we see that this case violates condition C. The same applies to the case when x is uneven.

Case ii.iii: Now it remains to consider when v and x are even. Suppose $v = 0^k 1 (0^{k+s} 1)^p 0^s$ for some $k, s, p \in \mathbb{N}$, and $x = 0^c 1 (0^{c+d} 1)^e 0^d$ for some $c, d, e \in \mathbb{N}$.

Case ii.iii.i: First of all we consider the case when $m(\text{red}, v) = 0$. Then $m(\text{red}, x) = 0$ since x precedes v in z . It follows that $m(\text{red}, vx) = 0$, which violates condition A.

Case ii.iii.ii: Now we turn to the case when $m(\text{red}, v) > 0$. By the same argument in Case i.iii, we get that $m(\text{green}, v) = 0$ and $k + s = 3$. Note that $p < n$, for otherwise the condition $m(\text{green}, v) = 0$ cannot be satisfied. Now we consider the following subcases:

Case ii.iii.ii.i: If $m(\text{red}, x) > 0$, then $m(\text{green}, x) = 0$ and $c + d = 3$. After selecting $i = \frac{n!}{2+p+e}$, we get that $z_i = uv^i wx^i y = w_4^{n!+n} w_3 w_2^{n!+n} w_3 w_3^{2(n!+n)} = (w_4^{n!+n} w_3 w_2^{n!+n}) (w_3 w_3^{2(n!+n)})$ is an abelian square, which violates condition C again.

Case ii.iii.ii.ii: If $m(\text{red}, x) = 0$ and $m(\text{green}, x) > 0$, then $m(\text{blue}, x) < 3$, for otherwise x cannot be even. There are again four subcases here.

1. The first subcase is $m(\text{blue}, x) = 2$. Then x is in the form $001r_1$, where r_1 is any word. We see that (a) $r_1 = \epsilon$ or (b) $r_1 = 0$, for otherwise x cannot be even. (a) If $r_1 = \epsilon$, then $x = 001$. We pick $i = 2$. It follows that $uv^2 wx^2 y = w_4^{n+2p+2} w_3^2 w_2^{n!+n} w_3 w_3^{2(n!+n)} \notin T$, which violates condition C. (b) If $r_1 = 0$, then $x = 0010$. We pick $i = 2$. It follows that $uv^2 wx^2 y = w_4^{n+2p+2} w_3 w_4 w_2^{n!+n} w_3 w_3^{2(n!+n)} \notin T$, which also violates condition C.
2. The second case is $m(\text{blue}, x) = 1$. Here is a picture.

$$z = w_4 w_4 \cdots w_4 10 \underbrace{0 w_2 \cdots \cdots w_2 w_3 \cdots w_3}_x$$

Thus (a) $x = 010$ or (b) $x = (01)^l$ for some $l > 0$. (a) If $x = 010$, then we pick $i = 2$. It follows that $uv^2 wx^2 y = w_4^{n+2p+2} w_3^2 w_2^{n!+n} w_3 w_3^{2(n!+n)} \notin T$. (b) Otherwise $x = (01)^l$ for some $l > 0$. After picking $i = \frac{n!}{1+p}$, we get

$$\begin{aligned} z_i &= w_4^{n!+n} w_3 w_2^{n!+n+\frac{ln!}{1+p}} w_3 w_3^{2(n!+n)} \\ &= (w_4^{n!+n} w_3 w_2^{n!+n}) w_2^{\frac{ln!}{1+p}} (w_3 w_3^{2(n!+n)}) \\ &= (w_4^{n!+n} w_3 w_2^{n!+n} w_2^{\frac{ln!}{2(1+p)}}) (w_2^{\frac{ln!}{2(1+p)}} w_3 w_3^{2(n!+n)}). \end{aligned}$$

Note that $\frac{ln!}{2(1+p)}$ is an integer since $n > 4$. Therefore z_i is an abelian square. Hence $z_i \notin T$, a contradiction.

3. The third case is $m(\text{blue}, x) = 0$ and $m(\text{green}, x) > 1$. Similarly x must be of the form $(01)^l$ or $(10)^l$ for some $l \in \mathbb{N}^+$, since x is even. We pick $i = \frac{n!}{1+p}$ and find the same result as in the second case.

4. The last case is exactly when $m(\text{blue}, x) = 0$ and $m(\text{green}, x) = 1$. Then x has to be the first or the last letter of the substring $w_2^{n!+n}$ of z , since x cannot be a single 0 or 1 (there are no successive 1's in elements of R). Moreover, we find that x cannot be the first letter of $w_2^{n!+n}$, since $m(\text{blue}, x) = 0$. It follows that x is the trailing 0 of $w_2^{n!+n}$. Under this circumstance, we find that $x = 0(100)^e 10$, since x is even. Now we select $i = \frac{n!}{1+p}$. It follows that

$$\begin{aligned}
z_i &= uv^i wx^i y \\
&= w_4^{n!+n} w_3 w_2^{n!+n} w_3 w_3^{2(n!+n) + \frac{(1+e)n!}{1+p}} \\
&= (w_4^{n!+n} w_3 w_2^{n!+n}) w_3^{\frac{(1+e)n!}{1+p}} (w_3 w_3^{2(n!+n)}) \\
&= (w_4^{n!+n} w_3 w_2^{n!+n} w_3^{\frac{(1+e)n!}{2(1+p)}}) (w_3^{\frac{(1+e)n!}{2(1+p)}} w_3 w_3^{2(n!+n)}).
\end{aligned}$$

Note that $\frac{(1+e)n!}{2(1+p)}$ is an integer since $n > 4$. Thus z_i is an abelian square. Therefore $z_i \notin T$, a contradiction.

Case ii.iii.ii.iii: The last possible subcase is when $m(\text{red}, x) = 0$ and $m(\text{green}, x) = 0$ and $m(\text{black}, x) > 0$. In this case, we get that $c + d = 2$. Now we pick $i = \frac{n!}{1+p}$. It follows that

$$\begin{aligned}
z_i &= uv^i wx^i y \\
&= w_4^{n!+n} w_3 w_2^{n!+n} w_3 w_3^{2(n!+n) + \frac{(1+e)n!}{1+p}} \\
&= (w_4^{n!+n} w_3 w_2^{n!+n}) w_3^{\frac{(1+e)n!}{1+p}} (w_3 w_3^{2(n!+n)}) \\
&= (w_4^{n!+n} w_3 w_2^{n!+n} w_3^{\frac{(1+e)n!}{2(1+p)}}) (w_3^{\frac{(1+e)n!}{2(1+p)}} w_3 w_3^{2(n!+n)}).
\end{aligned}$$

Note that $\frac{(1+e)n!}{2(1+p)}$ is an integer since $n > 4$. Thus z_i is an abelian square. Therefore $z_i \notin T$, a contradiction again.

With the above discussion, we claim that no decomposition of z can satisfy all three conditions simultaneously. Thus T is not context-free. $\square$

Proof. Combining Lemma 9 with the closure property, that the language obtained by intersecting a context-free language and a regular language is still context-free, we finally obtain this conclusion. $\square$

References

- [1] P. Erdős. Some unsolved problems. *Magyar Tud. Akad. Mat. Kutató Int. Közl.* **6** (1961), 221–264.